

Contents Volume 29

No. 1

- The initiation of development at fertilization (*Review*)
D. Epel (U.S.A.)

- Requirement for G protein activity at a specific time during embryonic chick myogenesis
R.E. Hausman, H. ElGendy and F. Craft (U.S.A.)

- Phalloidin inhibits cortical granule exocytosis and ooplasmic segregation in loach eggs
V.V. Ivanenkov, A.A. Minin and S.G. Ozerova (U.S.S.R.)

- Is ubiquitin involved in the dedifferentiation of higher plant cells?
E. Jamet, A. Durr, Y. Parmentier, M.-C. Criqui and J. Fleck (France)

- Characterization of MPF activation by okadaic acid in *Xenopus* oocyte
H. Rime, D. Huchon, C. Jessus, J. Goris, W. Merlevede and R. Ozon (France, Belgium)

- Effects of medium and substratum on ovine satellite cell attachment, proliferation and differentiation in vitro
M.V. Dodson, B.A. Mathison and B.D. Mathison (U.S.A.)

- The behaviour of cells from the distal tips of quail wing buds when grafted back into chick wings after micro-mass culture
C.P. Cottrill, A. Crawley and C. Tickle (U.K.)

No. 2

- Direct transdifferentiation: can cells change their phenotype without dividing?
W.A. Beresford (U.S.A.) (Review)

- Extracellular matrix production by embryonic epithelium cultured on type IV collagen. Deposition of a primary corneal stroma-like structure containing large irregular type I fibrils without type II collagen
F. Ruggiero, A. Barge, J.-L. Coll and R. Garrone (France)

- Leukocyte surface markers in *Rana catesbeiana*, identified using mouse monoclonal antibodies

- K. Sugiyama, A. Amenomori and K. Hatakeyama (Japan)*

- Uncoupling of muscle-specific protein expression in myocyte \times myoblast heterokaryons
W.E. Wright and V.K. Lin (U.S.A.)

- Electroporation as a new technique for producing transgenic fish
K. Inoue, S. Yamashita, J.-i. Hata, S. Kabeno, S. Asada, E. Nagahisa and T. Fujita (Japan)

- Regulation of oocyte number during oocyte differentiation in the lizard *Podarcis sicula*
P. Andreuccetti, C.M. Motta and S. Filosa (Italy)

- Maintenance of compaction and adherent-type junctions in mouse morula-stage embryos
I. Reima (Finland)

No. 3

Review

- Stem cell potential of ductular and periductular cells in the adult rat pancreas
M.S. Rao, A.V. Yeldandi and J.K. Reddy (U.S.A.)

Research Papers

- The silver-stained NOR and argentophilic nuclear proteins in early mouse embryogenesis: a cytological study
A.P. Dyban, E.L. Severova, O.V. Zatsepina and Y.S. Chentsou (U.S.S.R.)

- A mouse embryonic stem cell line showing pluripotency of differentiation in early embryos and ubiquitous β -galactosidase expression
H. Suemori, Y. Kadodawa, K. Goto, I. Araki, H. Kondoh and N. Nakatsuji (Japan)

- Cell lineage analyses of epithelia and blood vessels in chimeric mouse embryos by use of an embryonic stem cell line expressing the β -galactosidase gene
Y. Kadokawa, H. Suemori and N. Nakatsuji (Japan)

Development of cell surface activity and cell surface adhesiveness in early embryos of the newt, <i>Cynops pyrrhogaster</i> <i>S. Komazaki (Japan)</i>	195	14-kDa galactose-binding lectins possibly associated with an apolipoprotein <i>S.E. Zalik, W.J. Schneider and I.M. Ledsham (Canada)</i>	217
Structural modifications of the nuclear components during lizard oogenesis in relation to the differentiation of the follicular epithelium <i>C. Taddei and P. Andreuccetti (Italy)</i>	205	<i>Contents Vol. 29</i>	233
<i>Rapid Communication</i> The gastrulating chick blastoderm contains 16-kDa and		<i>Author Index</i>	235
		<i>Subject Index</i>	237

